

TOXIC
KEEP OUT OF REACH OF CHILDREN
ECOTOXIC

CURFEW[®]

FUNGICIDE

**A curative fungicide for control of downy mildew
in garlic and onions and late blight in potatoes**

READ THIS LABEL COMPLETELY BEFORE USE

ACTIVE INGREDIENT

Contains 450g/kg cymoxanil in the form of a
water dispersible granule

NET CONTENTS:
2.5kg



AGRECOVERY

Distributed by:
Etec Crop Solutions Limited
PO Box 51 584, Pakuranga, Auckland
Registered To: Agricultural Imports Direct NZ Ltd
Level 1, 101 Station Road,
Penrose, Auckland
New Zealand

Phone: 0800 100 325
www.etec.co.nz

Etec
CROP SOLUTIONS

HSNO CLASSIFICATIONS: 6.1D, 6.3B, 6.4A, 6.5B, 6.9B, 9.1B, 9.3C.

HARMFUL: May be harmful if swallowed, inhaled or absorbed through the skin. May cause sensitisation from prolonged skin contact. May cause eye irritation. Wear eye protection when mixing or applying. May cause organ damage from repeated oral exposure at high doses.

ECOTOXIC: Very Toxic to fish with long lasting effects. Avoid contamination of any water supply with chemical or empty container.

Do not apply **CURFEW** onto or near water. Harmful to terrestrial vertebrates.

PRECAUTIONS: Keep out of reach of children. Avoid skin contact and inhalation of spray mist or powder. Wash hands and exposed skin before meals and after work. Dispose of empty container safely. Keep away from fire or sparks.

CONTAINER DISPOSAL: Triple rinse container and add residue to spray tank. Submit clean empty container for recycling through Agrecovery. If this is not possible, burn if permitted and circumstances, especially wind direction, allow, otherwise bury in landfill.

HANDLING PRECAUTIONS: When mixing or applying, avoid contact with skin and eyes. Wear protective clothing, waterproof gloves and goggles. Do not eat or drink while using. Wash hands and face before meals and after work. Wash protective clothing after work.

STORAGE: Store in the original container, tightly closed and in a locked, dry cool well ventilated area away from feed, seeds and foodstuffs. Storage must be in accordance with the current version of NZS8409 Management of Agrichemicals and storage of certain quantities may require bunding and signage. See the Product Safety Card for this product for further information.

When stored appropriately this product should show no significant degradation for two years from the date of manufacture. Contact your supplier for further information about the use of any product that is older than this.

FIRST AID: If swallowed DO NOT induce vomiting. For advice, contact the National Poisons Centre 0800 Poison (0800 764 766) or a doctor immediately. If skin or hair contact occurs, remove contaminated clothing and flush skin and hair with running water. DO NOT SCRUB SKIN. If concentrate is splashed in the eyes, flush with running water immediately.

GENERAL DIRECTIONS: **CURFEW** is a contact and systemic fungicide that has curative activity on downy mildew and late blight, when applied immediately after spore germination. It also acts as a short term protectant.

Resistance Management: Always apply **CURFEW** in mixture with mancozeb fungicide.

See 'Directions for Use' for recommendations. Do not use **CURFEW** more than three times in succession and do not apply more than 5 times per season.

RAINFALL: Do not spray if rainfall is expected within 1 hour of spraying.

COMPATIBILITY: **CURFEW** is compatible with most insecticides, but it is not compatible with Bordeaux mixture, lime-sulphur or other alkaline products.

MIXING: Add the required amount of **CURFEW** directly into the partly filled spray tank whilst agitating. Ensure Curfew is mixed thoroughly before adding any other pesticide to the spray tank. Agitate during use and before using after a stoppage.

Use all spray as soon as possible after mixing and do not allow to stand for an extended period, such as overnight.

APPLICATION: DIRECTIONS FOR USE: As indicated below, **CURFEW** must be used in combination with an appropriate protectant fungicide, and is intended to be used as part of a spray programme with alternative fungicides.

CROP	DISEASE	REMARKS
Garlic, Onions	Downy Mildew	Apply 250-350g/ha as a tankmix partner with mancozeb at the recommended label rates. Begin spraying at the 2 true leaf stage and continue at 7-10 day intervals. Apply CURFEW immediately after an infection period (but before disease symptoms occur). Do not use more than three times in succession and do not apply more than 5 times per season.
Potatoes	Late Blight	Apply 220-250g/ha as a tankmix partner with mancozeb at the recommended label rates at 7-10 day intervals. Use CURFEW , applying immediately after an infection period (but before disease symptoms occur) and use alternative fungicides at other times. Do not use more than three times in succession and do not apply more than 5 times per season. Use the higher rate of CURFEW under extreme infection conditions (Warm with high humidity).


CONDITIONS OF SUPPLY: All goods supplied by us are produced under rigid quality control procedures which assure the purchaser that at the time of sale they are physically suitable for use in the manner stated. However, to the extent permitted by law, buyers and/or users of the goods accept that neither the manufacturer, Agricultural Imports Direct NZ Ltd as distributor, nor any other distributor have any liability or responsibility whatsoever for any loss, damage or injury whether in contract or tort, whether direct, indirect or consequential howsoever arising in connection with the supply of these goods.

Registered pursuant to the ACVM Act 1997, No. P8010
See www.nzfsa.govt.nz/acvm/ for registration conditions
Approved pursuant to the HSNO Act 1996, No. HSR100090
See www.ermanz.govt.nz for approval controls

Batch No:

DOM: See sticker on box

UN No. 3077
ENVIRONMENTALLY
HAZARDOUS SUBSTANCE,
SOLID, N.O.S (CYMOXANIL)
Class: 9
Packing Group: III



EL346

24 Hour Emergency Response
0800CHEM CALL
0800 243 622



9 421022 422563

dieline

pms 512

pms646

black

pm369

pm375

FRONT Label 140x140

BACK Label