

**FLAMMABLE LIQUID**  
**KEEP AWAY FROM SOURCES OF IGNITION**

# Chloro IPC

*A herbicide for use post-emergence in onions, pre- and post-emergence in lettuce and pre-emergence in carrots, leeks and parsnips.*

**Read the Label Carefully before use**

**Active Ingredient:**

400 g/litre chlorpropham in the form of an emulsifiable concentrate.  
Also contains 486 g/litre hydrocarbon liquid.

**HERBICIDE**



**AGRECOVERY**

Registrant:

Elliott Technologies Limited  
Box 838 Pukekohe  
New Zealand

**Phone: 0800 100 325**

**www.elliott-technologies.co.nz**

**NET CONTENTS:**  
**10 Litres**



170mm deep

190mm wide

**HARMFUL** Keep out of reach of children. **ECOTOXIC**

**HSNO Class: 3.1C, 6.1E, 6.3B, 6.4A, 6.9B, 9.1B, 9.2A**

**Approved Handler and Tracking apply**

This product must be under the personal control of an approved handler or secured Location and movement of this product must be recorded at each stage of its life cycle

**ENVIRONMENTAL WARNING: TOXIC TO AQUATIC ORGANISMS.**  
DO NOT apply to bodies of water. DO NOT contaminate sewers, drains, dams, creeks or any other waterways with product or empty container.

**VERY TOXIC IN THE SOIL ENVIRONMENT.** Very toxic to some plant species – selective herbicide.

**HEALTH WARNING: HARMFUL** – may be harmful if swallowed, inhaled or absorbed through the skin. May cause skin and eye irritation. Avoid contact with skin and eyes. Repeated oral exposure at high dose is presumed to cause damage to the blood system.

**PRODUCT DISPOSAL:** Ensure unwanted product is used by another operator strictly in accordance with label directions.

**CONTAINER DISPOSAL:** Triple rinse container and add residue to spray tank. Submit clean empty container for recycling through Agrecovery. If this is not possible, burn if permitted and circumstances, especially wind direction allow, otherwise bury in landfill

**HANDLING PRECAUTIONS:** When mixing or applying, wear protective clothing, gloves and goggles. Do not eat drink or smoke while using. Wash hands and face before meals and after work. Wash protective clothing after work. Flush spray equipment thoroughly with clean water after use.

**STORAGE:** Store in original container tightly closed and in a locked, dry, cool, well ventilated area, away from feed, seeds and foodstuffs. Do not store or use near sources of ignition. This product should be stored in accordance with NZS 8409 Management of Agrichemicals and storage of certain quantities may require bunding and signage. See the HAZNOTE for further information.

When stored appropriately this product should show no significant degradation for 2 years. Contact your supplier for further information about the use of any product that is older than this.

**First Aid:** If swallowed do NOT induce vomiting. For advice, contact the **National Poisons Centre 0800 POISON (0800 764 766)** or a doctor immediately. If skin or hair contact occurs, remove contaminated clothing and flush skin and hair with running water.

**PRODUCT INFORMATION**

**CHLORO IPC** is mainly a soil acting residual herbicide with generally weak contact action. It is taken up by the emerging shoot and roots of weeds and affects the growing point of certain plants post-emergence. Its mode of action is disruption of cell division, transpiration and respiration and also protein synthesis. Rain or irrigation is required shortly after application for residual control of weeds.

**WEEDS CONTROLLED (Seedlings only)**

Black nightshade	Chickweed	Docks	Fathen
Foxtail	Fumitory	Paspalum	Annual Poa
Italian ryegrass	Speedwells	Spurrey	Willow Weed

**DIRECTIONS FOR USE**

**Mixing:** Half fill the spray tank with water and, with agitator running, add **CHLORO IPC** and continue filling the tank.

If tank mixing with other products, follow mixing instructions on the other labels.

**Application:** Apply to a firm moist seed bed in 200 - 400 litres water per hectare.

**Recommendations:**

**Carrots, leeks, parsnips - Pre-emergence** - use 6 - 8 litres / ha immediately after sowing.

**Lettuce - Pre-emergent** - Apply 3 - 8 litres at time of drilling.

**- Pre-transplanting** - Cultivate the soil into a firm seed bed and apply 6 - 11 litres per hectare prior to transplanting. Do not cultivate soil after application. Transplanting should be done to keep soil disturbance to a minimum and treated soil should not be incorporated into the root zone of the lettuce plants.

**Onions - Post-emergence** - Apply 1 - 2 litres per hectare at the flag stage of the onion, in combination with Chloronion™ at 1 - 2 litres/ ha plus Peptoil™ Crop Oil Concentrate at 1 litre/ ha. Repeat this mixture 2 - 3 weeks later.

- NOTES:**
1. Best results will be obtained if application is followed by 15 mm of rain or irrigation.
  2. Use the lower rates on silty or sandy soils.
  3. Excessive rainfall may wash the product too deeply into the soil and retard crop growth, particularly on light or sandy soils.

**Exclusion of Liability: Please Read**

To the extent permitted by law, buyers and/or users of the goods accept that neither the manufacturer, Elliott Technologies Limited as distributor, nor any other distributor have any liability or responsibility whatsoever for any loss, damage or injury whether in contract or tort, whether direct, indirect or consequential howsoever arising in connection with the supply of these goods.

See [www.nzfsa.govt.nz/acvm/](http://www.nzfsa.govt.nz/acvm/) for registration conditions.

**24 Hour emergency response**  
**0800CHEM CALL**  
**0800 243 622**

® - Registered Trademark of Elliott Technologies Limited  
TM<sup>1</sup> - Trade Mark of Drexel Chemical Company, USA

Batch No.  
DOM

Registered pursuant to the ACVM Act 1997, No. P5027  
HSNO Approval Code HSR000823



EL244

9 421022 420286

170mm deep

190mm wide

Dieline

PMS 363

PMS 485

Black