

Ovation[®] 50WDG

INSECTICIDE

- 500 g/kg buprofezin as a water dispersible granule
- Pack size: 1kg
- Insect Growth Regulator that interferes with the young developmental stages of certain insects. Has no effect on adults
- Activity is gradual
- Residual activity
- IPM friendly
- Low toxicity and low environmental impact

Benefits:

- ✓ Very Cost Effective
- ✓ Soft on beneficial Insects including bumblebees, parasitoid wasps and Encarsia sp.
- ✓ Mealy bug
- ✓ Scale
- ✓ Whitefly

- ✓ Grapes
- ✓ Apples
- ✓ Stonefruit
- ✓ Persimmons & Tamarillos
- ✓ Certain Glasshouse Vegetables



Ovation® 50WDG

GETTING THE BEST OUT OF OVATION:

- Coverage is critical – Use Bond Xtra® for medium volume spraying and Du-Wett® or Du-Wett Rainmaster® for low volume spraying.
- Use timely applications based on crop monitoring and targeted to the correct pest stage.

Use Recommendations

Use 25g / 100 L on all crops. Note recommended water rates for crops.

The rate of 25g / 100 litres is for high volume application to the point of run-off. For concentrate application, adjust dilution rates accordingly.

Applications in all crops should ensure that not less than 250g/ha are applied.

CROP	PEST	REMARKS
Citrus	Scales	For dilute spraying use 2500-3000 L/ha or more if required to obtain complete coverage. Treat when scale first appears and repeat 10-14 days later. Where populations persist, re-apply at intervals not less than one month. For season long programmes, alternate Ovation 50WDG with other scale materials. Adjust water rate for low volume spraying with Du-Wett but use the same rate of product per hectare as dilute.
Grapes	Mealy bug	For dilute spraying use 1000 L/ha of water (250g /ha minimum dose rate). Adjust water rate for low volume spraying. Apply the first spray in mixture with 1% spraying oil at the first sign of bud movement. Make a second application without oil 14-21 days later when shoots are 10 – 20 mm long. Do not apply after the start of flowering.
Kiwifruit	Greedy scale & oleander scale	Make one application between green tip and opening of the first female flowers. Where more than one preflower insecticide is required, use Ovation 50WDG for only one of these.
Persimmon	Mealy bug	Apply two applications at 10 – 14 days between green tip and bloom. Use sufficient water to obtain complete plant coverage.
Pipfruit	Mealy bug	Apply two applications at 10 – 14 days between green tip and the commencement of pink. Use sufficient water to obtain complete plant coverage. Consider concentrate spraying with Du-Wett, Oil and Oilmate® for improved efficacy.
Tamarillo	Whitefly	Apply after pruning and before fruit set. Re-apply at no less than 28 day intervals when white fly adults exceed 20 per leaf. Alternate with other whitefly materials to control adults.
Glasshouse Crops-Cucumbers, eggplant, melons, peppers, pepino, tomatoes & zucchini.	Whitefly	Apply as a drenching spray at the first appearance of whitefly and repeat if necessary at intervals of not less than one month. Apply no more than 2 applications of Ovation 50WDG per crop or season.

Warning: DO NOT USE ON ANY ORNAMENTAL CROP INCLUDING THOSE GROWN FOR CUT FLOWERS, FOLIAGE AND NURSERY PLANTS

©Ovation is a registered trademark of Etec Crop Solutions Ltd
Ovation is registered under the ACVM Act 1997 No. P5861

Distributed by:

Etec Crop Solutions Ltd. PO Box 51-584 Pakuranga, Auckland, New Zealand

NZ Freephone: 0800 100 325

www.etec.co.nz

