

KEEP OUT OF REACH OF CHILDREN

PROMETRYNE 500 SC

A selective herbicide for weed control in
carrots, parsnips, onions, peas and potatoes

Read the Label Carefully before use

Active Ingredient:
500 g/litre prometryne in the form
of a suspension concentrate.

NET CONTENTS
5 Litres

HERBICIDE
AGRECOVERY

Registrant:
Elliott Technologies Limited
Box 838 Pukekohe
New Zealand
Phone: 0800 100 325
www.elliott-technologies.co.nz



165mm deep

160mm wide

SHAKE CONTAINER WELL BEFORE USING

ECOTOXIC

HSNO Class: 6.1E, 6.9B, 9.1A, 9.2A

Approved Handler applies

This product must be under the personal control of an approved handler or secured.

ENVIRONMENTAL WARNING: VERY TOXIC TO AQUATIC ORGANISMS. DO NOT contaminate sewers, drains, dams, creeks or any other waterways with product or empty container. **VERY TOXIC TO THE SOIL ENVIRONMENT.** Very toxic to some plant species – selective herbicide.

HEALTH WARNING: HARMFUL – may be harmful if swallowed, inhaled or absorbed through the skin. Presumed to cause damage to the liver, kidneys and bone marrow through repeated oral exposure at high doses.

PRODUCT DISPOSAL: Ensure unwanted product is used by another operator strictly in accordance with label directions.

CONTAINER DISPOSAL: Triple rinse container and add residue to spray tank. Submit clean empty container for recycling through Agrecovery. If this is not possible, burn if permitted and circumstances, especially wind direction allow, otherwise bury in landfill

HANDLING PRECAUTIONS: When mixing or applying, wear protective clothing, gloves and goggles. Do not eat drink or smoke while using. Wash hands and face before meals and after work. Wash protective clothing after work.

STORAGE: Store in original container tightly closed and in a locked, dry, cool, well ventilated area, avoiding temperatures below -5° and above 30°C, away from feed, seeds and foodstuffs. This product is classified under HSNO as 9.1A, and storage of certain quantities may require bunding and signage. See the HAZNOTE for further information.

When stored appropriately this product should show no significant degradation for 2 years. Contact your supplier for further information about the use of any product that is older than this.

First Aid: If swallowed do NOT induce vomiting. For advice, contact the National Poisons Centre 0800 POISON (0800 764 766) or a doctor immediately. If skin or hair contact occurs, remove contaminated clothing and flush skin and hair with running water.

EXCLUSION OF LIABILITY: PLEASE READ

To the extent permitted by law, buyers and/or users of the goods accept that neither the manufacturer, Elliott Technologies Limited as distributor, nor any other distributor have any liability or responsibility whatsoever for any loss, damage or injury whether in contract or tort, whether direct, indirect or consequential howsoever arising in connection with the supply of these goods.

See www.nzfsa.govt.nz/acvm/ for registration conditions.

Registered pursuant to the ACVM Act 1997, No. P5654

HSNO Approval Code HSR000558

24 Hour emergency response

0800 CHEM CALL

0800 243 622

® Registered trademark of Elliott Technologies Limited

PRODUCT INFORMATION

Prometryne 500SC is an easy-to-use flowable formulation. Prometryne 500SC is a selective contact and residual herbicide providing a quick knockdown with short term residual activity on a wide range of broad leaf weeds and annual grasses. Prometryne 500SC gives best results when applied to small actively growing weeds in moist soil, free from clods.

Rainfall within 6 hours of application may reduce foliar knockdown. Rainfall is required however for residual control of germinating seedlings

Weeds Controlled:

Annual poa	Chickweed	Cornbind	Fumitory
Mayweeds *	Speedwell *	Spurrey	Twin cress *
Wild turnip	Willow weed	Wireweed *	

*These weeds may not be fully controlled when large or when low use rates are used.

DIRECTIONS FOR USE

MIXING: Always shake the container a few times before use. Pour the required amount of Prometryne 500SC into the partly filled spray tank with agitation running. Complete filling of spray tank. Maintain agitation before and during spraying and before resuming spraying after a stoppage.

RECOMMENDATIONS: Apply Prometryne 500SC in 200-500 litres of water per hectare. Use the higher water rate to improve selectivity on onions and peas but avoid misting. Do not add a surfactant as this may increase the risk of crop damage. Do not use strainers finer than 300 microns. Flush equipment with clean water after use.

RATES AND TIMING OF APPLICATION:

CROP	TIMING	RATE/HECTARE	COMMENTS
Carrots Parsnips	minimum 2 true leaves	1.5-2 litres	Apply while weeds are still small
Onions	minimum 2 true leaves	650 – 750 ml	Repeat applications can be made if further germinations occur. Do not spray within 7 days of fungicide or insecticide spray as damage may occur. Do not spray when onions are bulbing.
Potatoes	maximum 5% emergence	2 litres	Apply after final moulding while weeds are still small.
Peas	5 – 8cm high	1 litre – 1.5 litres	Apply to disease free crops not more than 10cm high. Allow 2 days drying after heavy rain before spraying. NOTE: as peas are a tender crop which depend on the waxy leaf surface for selectivity, use a crystal violet dye retention test to confirm the presence of sufficient leaf wax before spray application.

Compatibility:

- Do not use with oils or emulsifiable concentrates unless stated on label.
- Do not mix with another herbicide unless weather and crop conditions are ideal for use of both products according to label recommendations.

Batch No.

DOM

EL245



9 421022 420880

165mm deep

Dieline

PMS 363

PMS 485

Black

160mm wide