

KEEP OUT OF REACH OF CHILDREN

super sprout
Stop™

Growth regulator for the prevention of
sprouting of Onions and Potatoes

Read the Label Carefully before use

Active Ingredient: 270 g/litre maleic hydrazide
as the potassium salt in the form of a soluble concentrate.

SPROUT INHIBITOR



NET CONTENTS:
10 Litres

Registered By:
Elliott Technologies Limited
P.O.Box 838, Pukekohe, New Zealand.
Phone: 0800 100 325 www.elliott-technologies.co.nz



super sprout
Stop™

ECOTOXIC

HSNO Class: 9.2A
Approved Handler

This product must be under the control of an approved handler or secured.

ENVIRONMENTAL WARNING: VERY TOXIC TO THE SOIL ENVIRONMENT

PRODUCT DISPOSAL: Ensure unwanted product is used by another operator strictly in accordance with label directions.

CONTAINER DISPOSAL: Triple rinse container and add residue to spray tank. Burn if permitted and circumstances, especially wind direction allow, otherwise bury in landfill.

HANDLING PRECAUTIONS: When mixing or applying, wear protective clothing, gloves and goggles. Do not eat or drink while using. Wash hands and face before meals and after work. Wash protective clothing after work. Flush spray equipment thoroughly after use.

STORAGE: Store in original container tightly closed and in a locked, dry, cool, well ventilated area, away from foodstuffs Storage must be generally in accordance with NZS8409 Management of Agrichemicals. See the HAZNOTE for further information.

When stored appropriately this product should show no significant degradation for 2 years from date of manufacture. Contact your supplier for further information about the use of any product that is older than this.

SPILLAGE: Wear appropriate protective clothing and prevent material from entering waterways. Absorb spills with inert material and place in waste containers. Sweep up solids without creating dust. Wash area with water and absorb with further inert material. Dispose of waste safely (such as to a suitable landfill).

FIRST AID: If swallowed **NOT** induce vomiting. For advice, contact the National Poisons Centre **0800 poison (0800 764 766)** or a doctor immediately. If skin or hair contact occurs, remove contaminated clothing and flush skin and hair with running water.

PRODUCT INFORMATION

Super Sprout Stop is a liquid formulation of the potassium salt of maleic hydrazide. It is absorbed by the green foliage of the plant and is translocated by the plants fluids to the growth areas where cell division occurs. **Super Sprout Stop** inhibits cell division and prevents onions and potatoes from sprouting.

WARNING: There are an increasing number of new varieties of both onions and potatoes becoming available to growers. Some of these varieties may require higher or lower than normal rates of **Super Sprout Stop** to achieve the expected result. Until experience has been obtained with a previously untested variety, treat only a small area in the first instance.

DIRECTIONS FOR USE

MIXING: **Super Sprout Stop MIXES EASILY WITH WATER.** Pour the required quantity into the partly filled spray tank with agitation running and complete filling of the spray tank.

APPLICATION: **Super Sprout Stop** may be applied with any conventional spray equipment either by ground or air.

NOTE: DO NOT SPRAY IF RAIN IS LIKELY TO FALL WITHIN 24 HOURS AS THE EFFECTIVENESS MAY BE REDUCED.

RECOMMENDATIONS:

ONIONS: Apply 8 – 9 litres of **Super Sprout Stop** per hectare in 450 – 900 litres of water when bulbs are fully matured, but still have 5 – 8 green leaves. Best results are obtained when at least 50% of tops are still green. This is usually 10 – 14 days before harvest time. Add a non-ionic surfactant at the low label rate.

NOTE: Do not spray onions earlier than 2 weeks before maturity. This can result in spongy bulbs and deformed onions.

POTATOES: Apply 12 – 14 litres **Super Sprout Stop** per hectare in 450 – 900 litres of water. Check your field to see if you have a good set of potatoes. Apply **Super Sprout Stop** when most of the tubers are approximately 2.5 – 3.5 cm (1 – 1.5 inches) in diameter. This usually occurs approximately 1 week after blossoms fall and a few lower leaves show a yellow colour. Add a non-ionic surfactant at the low label rate.

Notes: 1. Do not spray potatoes earlier than recommended as yield may be reduced.

2. Do not treat potatoes grown for seed production.

EXCLUSION OF LIABILITY: PLEASE READ

To the extent permitted by law, buyers and/or users of the goods accept that neither the manufacturer, Elliott Technologies Ltd as distributor, nor any other distributor have any liability or responsibility whatsoever for any loss, damage or injury whether in contract or tort, whether direct, indirect or consequential howsoever arising in connection with the supply of these goods.

Registered pursuant to ACVM Act 1997, No.P5860. HSNO Approval Code HSR000798. See www.nzfsa.govt.nz/acvm/ for registration conditions.

™ - is a registered trademark of Drexel Chemical Company, USA.

® - is a registered trademark of Elliott Technologies Ltd

Batch No.

DOM

24 Hour Emergency Response
0800 CHEM CALL
0800 243 622



9 421022 420965

EL217



PMS 021



PMS 485



BLACK

150W X 180H